


CLAIMS

1. A compound of formula (I)



wherein

- A is a bivalent radical selected from $-C(O)-$, $-C(O)NH-$, $-NHC(O)-$, $-N(R^7)-CH_2-$, $-CH_2-N(R^7)-$, $-CH(NR^8R^9)-$ and $-C(=NR^{10})-$;

R^1 is $-O(CH_2)_dXR^{11}$;


R^2 is hydrogen or a hydroxyl protecting group;

R^3 is hydrogen, C_{1-4} alkyl, or C_{3-6} alkenyl optionally substituted by 9 to 10 membered fused bicyclic heteroaryl;

- R^4 is hydroxy, C_{3-6} alkenyloxy optionally substituted by 9 to 10 membered fused bicyclic heteroaryl, or C_{1-6} alkoxy optionally substituted by C_{1-6} alkoxy or $-O(CH_2)_eNR^7R^{12}$,

R^5 is hydroxy, or

R^4 and R^5 taken together with the intervening atoms form a cyclic group having the following structure:



wherein Y is a bivalent radical selected from $-CH_2-$, $-CH(CN)-$, $-O-$, $-N(R^{13})-$ and $-CH(SR^{13})-$;

R^6 is hydrogen or fluorine;


R^7 is hydrogen or C_{1-6} alkyl;

- R^8 and R^9 are each independently hydrogen, C_{1-6} alkyl, $-C(=NR^{10})NR^{14}R^{15}$ or $-C(O)R^{14}$, or

R^8 and R^9 together form $=CH(CR^{14}R^{15})_f$ aryl, $=CH(CR^{14}R^{15})_f$ heterocyclyl, $=CR^{14}R^{15}$ or $=C(R^{14})C(O)OR^{14}$, wherein the alkyl, aryl and heterocyclyl groups are optionally substituted by up to three groups independently selected from R^{16} ;


R^{10} is $-OR^{17}$, C_{1-6} alkyl, $-(CH_2)_g$ aryl, $-(CH_2)_g$ heterocyclyl or $-(CH_2)_hO(CH_2)_iOR^7$,
 5 wherein each R^{10} group is optionally substituted by up to three groups independently selected from R^{16} ;

R^{11} is a heterocyclic group having the following structure:



10

or



R^{12} is hydrogen or C_{1-6} alkyl;

15 R^{13} is hydrogen or C_{1-4} alkyl optionally substituted by a group selected from optionally substituted phenyl, optionally substituted 5 or 6 membered heteroaryl and optionally substituted 9 to 10 membered fused bicyclic heteroaryl;

R^{14} and R^{15} are each independently hydrogen or C_{1-6} alkyl;



20 R^{16} is halogen, cyano, nitro, trifluoromethyl, azido, $-C(O)R^{21}$, $-C(O)OR^{21}$, $-OC(O)R^{21}$, $-OC(O)OR^{21}$, $-NR^{22}C(O)R^{23}$, $-C(O)NR^{22}R^{23}$, $-NR^{22}R^{23}$, hydroxy, C_{1-6} alkyl, $-S(O)_kC_{1-6}$ alkyl, C_{1-6} alkoxy, $-(CH_2)_m$ aryl or $-(CH_2)_m$ heteroaryl, wherein the alkoxy group is optionally substituted by up to three groups independently selected from $-NR^{14}R^{15}$, halogen and $-OR^{14}$, and the aryl and heteroaryl groups are optionally substituted by up to five groups independently selected from halogen, cyano, nitro, trifluoromethyl, azido, $-C(O)R^{24}$, $-C(O)OR^{24}$, $-OC(O)OR^{24}$, $-NR^{25}C(O)R^{26}$, $-C(O)NR^{25}R^{26}$, $-NR^{25}R^{26}$, hydroxy, C_{1-6} alkyl and C_{1-6} alkoxy;

25 R^{17} is hydrogen, C_{1-6} alkyl, C_{3-7} cycloalkyl, C_{3-6} alkenyl or a 5 or 6 membered heterocyclic group, wherein the alkyl, cycloalkyl, alkenyl and heterocyclic groups are optionally substituted by up to three substituents independently selected from optionally substituted 5 or 6 membered heterocyclic group, optionally substituted 5 or 6 membered heteroaryl, $-OR^{27}$, $-S(O)_nR^{27}$, $-NR^{27}R^{28}$, $-CONR^{27}R^{28}$, halogen and cyano;


30 R^{18} is hydrogen, $-C(O)OR^{29}$, $-C(O)NHR^{29}$, $-C(O)CH_2NO_2$ or $-C(O)CH_2SO_2R^7$;

- R¹⁹ is hydrogen, C₁₋₄alkyl optionally substituted by hydroxy or C₁₋₄alkoxy, C₃₋₇cycloalkyl, or optionally substituted phenyl or benzyl;
- R²⁰ is halogen, C₁₋₄alkyl, C₁₋₄thioalkyl, C₁₋₄alkoxy, -NH₂, -NH(C₁₋₄alkyl) or -N(C₁₋₄alkyl)₂;
- 5 R²¹ is hydrogen, C₁₋₁₀alkyl, -(CH₂)_paryl or -(CH₂)_pheteroaryl;
- R²² and R²³ are each independently hydrogen, -OR¹⁴, C₁₋₆alkyl, -(CH₂)_qaryl or -(CH₂)_qheterocyclyl;
- R²⁴ is hydrogen, C₁₋₁₀alkyl, -(CH₂)_raryl or -(CH₂)_rheteroaryl;
- R²⁵ and R²⁶ are each independently hydrogen, -OR¹⁴, C₁₋₆alkyl, -(CH₂)_saryl or -(CH₂)_sheterocyclyl;
- 10 R²⁷ and R²⁸ are each independently hydrogen, C₁₋₄alkyl or C₁₋₄alkoxyC₁₋₄alkyl;
- R²⁹ is hydrogen,
 C₁₋₆alkyl optionally substituted by up to three groups independently selected from
 halogen, cyano, C₁₋₄alkoxy optionally substituted by phenyl or C₁₋₄alkoxy, -
 15 C(O)C₁₋₆alkyl, -C(O)OC₁₋₆alkyl, -OC(O)C₁₋₆alkyl, -OC(O)OC₁₋₆alkyl, -
 C(O)NR³²R³³, -NR³²R³³ and phenyl optionally substituted by nitro or -C(O)OC₁₋₆alkyl,
 -(CH₂)_wC₃₋₇cycloalkyl,
 -(CH₂)_wheterocyclyl,
 20 -(CH₂)_wheteroaryl,
 -(CH₂)_waryl,
 C₃₋₆alkenyl, or
 C₃₋₆alkynyl;
- R³⁰ is hydrogen, C₁₋₄alkyl, C₃₋₇cycloalkyl, optionally substituted phenyl or benzyl, acetyl
 25 or benzoyl;
- R³¹ is hydrogen or R²⁰, or R³¹ and R¹⁹ are linked to form the bivalent radical -O(CH₂)₂- or -(CH₂)_t;
- R³² and R³³ are each independently hydrogen or C₁₋₆alkyl optionally substituted by phenyl or -C(O)OC₁₋₆alkyl, or
- 30 R³² and R³³, together with the nitrogen atom to which they are bound, form a 5 or 6 membered heterocyclic group optionally containing one additional heteroatom selected from oxygen, nitrogen and sulfur;
- X is -U(CH₂)_vB-, -U(CH₂)_v- or a group selected from:


35



and




- U and B are independently a divalent radical selected from $-N(R^{30})-$, $-O-$, $-S(O)_Z-$, $-N(R^{30})C(O)-$, $-C(O)N(R^{30})-$ and $-N[C(O)R^{30}]-$;
- 5 W is $-C(R^{31})-$ or a nitrogen atom;
d is an integer from 2 to 6;
e is an integer from 2 to 4;
f, g, h, m, p, q, r, s and w are each independently integers from 0 to 4;
i is an integer from 1 to 6;
- 10 j, k, n and z are each independently integers from 0 to 2;
t is 2 or 3;
v is an integer from 1 to 8;
or a pharmaceutically acceptable derivative thereof.
- 15 2. A compound according to claim 1 wherein A is $-C(O)-$ or $-N(R^7)-CH_2-$.
3. A compound according to claim 1 or claim 2 wherein X is $-U(CH_2)_vB-$ or $-U(CH_2)_v$.
4. A compound according to any one of the preceding claims wherein d is 2 or 3.
- 20 5. A compound according to any one of the preceding claims wherein R^{11} is a heterocyclic group of the following formula:




25

or




wherein the heterocyclic is linked in the 6 or 7 position and j, R¹⁸, R¹⁹ and R²⁰ are as defined in claim 1;

5 a heterocyclic group of the following formula:



wherein the heterocyclic is linked in the (ii) or (iii) position, W is -C(R³¹)- and R³¹ and R¹⁹ are linked to form the bivalent radical -(CH₂)_t- as defined in claim 1, and j, R¹⁸, R¹⁹ and R²⁰ are as defined in claim 1; or
10 a heterocyclic group of the following formula:



15 wherein the heterocyclic is linked in the 7 or 8 position and j, R¹⁸, R¹⁹ and R²⁰ are as defined in claim 1.

6. A compound according to claim 1 as defined in any one of Examples 1 to 42, or a pharmaceutically acceptable derivative thereof.

20

7. A compound selected from:

4"-O-(2-[[2-(3-carboxy-1-cyclopropyl-6-fluoro-4-oxo-1,4-dihydro-quinolin-7-ylamino)-ethyl]-methylamino]-ethyl)-6-O-methyl-erythromycin A 11,12-carbonate;

4"-O-(3-[[2-(3-carboxy-1-cyclopropyl-6-fluoro-4-oxo-1,4-dihydro-quinolin-7-ylamino)ethyl]-methylamino]-propyl)-6-O-methyl-erythromycin A 11,12-carbonate;


25



4"-O-{3-[2-(2-carboxy-1-oxo-6,7-dihydro-1*H*,5*H*-pyrido[3,2,1-*ij*]quinoline-9-yloxy)-ethylamino]-propyl}-6-O-methyl-erythromycin A 11,12-carbonate;

- 4"-O-(3-[[3-(3-carboxy-1-ethyl-4-oxo-1,4-dihydro-quinolin-6-yl)propyl]-methylamino]-propyl)-6-O-methyl-erythromycin A 11,12-carbonate;
 4"-O-(3-[[2-(3-carboxy-1-ethyl-6-fluoro-4-oxo-1,4-dihydro-[1,8]naphthyridin-7-ylamino)ethyl]-methylamino]-propyl)-6-O-methyl-erythromycin A 11,12-carbonate;
 5 4"-O-{2-[2-(3-carboxy-1-ethyl-6-fluoro-4-oxo-1,4-dihydro-[1,8]naphthyridin-7-ylamino)ethyl]-methylamino}-ethyl }-6-O-methyl-erythromycin A;
 4"-O-{3-[[3-(3-carboxy-1-ethyl-4-oxo-1,4-dihydro-quinolin-6-yl)-propyl]-methylamino]-propyl}-6-O-methyl-11-desoxy-11-(R)-amino-erythromycin A 11,12-carbamate;
 4"-O-{3-[[2-(3-carboxy-1-ethyl-4-oxo-1,4-dihydro-quinolin-6-ylsulfanyl)-ethyl]-methylamino]-propyl}-6-O-methyl-11-desoxy-11-(R)-amino-erythromycin A 11,12-carbamate;
 10 4"-O-{3-[2-(3-carboxy-7-chloro-1-cyclopropyl-4-oxo-1,4-dihydro-quinolin-6-ylamino)-ethylcarbamoyl]-propyl}-azithromycin;
 4"-O-{2-[2-(3-carboxy-6-fluoro-1-cyclopropyl-4-oxo-1,4-dihydro-quinolin-7-ylamino)-ethylamino]-ethyl}-azithromycin 11,12-cyclic carbonate;
 15 4"-O-{2-[2-(3-carboxy-7-chloro-1-cyclopropyl-4-oxo-1,4-dihydro-quinolin-6-ylamino)-ethylamino]-ethyl}-azithromycin; and
 4"-O-{2-[2-(3-carboxy-6-fluoro-1-cyclopropyl-4-oxo-1,4-dihydro-quinolin-7-ylamino)-ethylamino]-ethyl}-azithromycin;
 20 or a pharmaceutically acceptable derivative thereof.

8. A process for the preparation of a compound as claimed in claim 1 which comprises:


- 25 a) reacting a compound of formula (II)





5 with a suitable amine (IIIa) or (IIIb), wherein B^a and R^{11a} are B and R^{11} as defined in claim 1 or groups convertible to B and R^{11} ;

b) reacting a compound of formula (V)



(V)

10


with a compound of formula X^aR^{11a} (IV), wherein R^{11a} is R^{11} as defined in claim 1 or a group convertible to R^{11} and X^a is $-\text{U}(\text{CH}_2)_V-$ or $-\text{U}(\text{CH}_2)_V\text{B}-$, or a group convertible to $-\text{U}(\text{CH}_2)_V-$ or $-\text{U}(\text{CH}_2)_V\text{B}-$, in which U is a group selected from $-\text{N}(\text{R}^{30})-$ and $-\text{S}-$, and L is suitable leaving group, to produce a compound of formula (I) wherein U is a group selected from $-\text{N}(\text{R}^{30})-$ and $-\text{S}-$;

15

c) converting one compound of formula (I) into another compound of formula (I);

d) where U is $-\text{O}-$, reacting a compound of formula (VII)


20



(VII)

with a suitable compound of formula X^aR^{11a} in the presence of a catalyst; or

- 5 e) where U is $-C(O)N(R^{30})-$, reacting a compound of formula (VIII)



with a suitable amine compound,

10

and thereafter, if required, subjecting the resulting compound to one or more of the following operations:

- i) removal of the protecting group R²,
 ii) conversion of X^aR^{11a} to XR^{11} ,
 15 iii) conversion of B^aR^{11a} to R^{11} ,
 iv) conversion of R^{11a} to R^{11} ,

and

v) conversion of the resultant compound of formula (I) into a pharmaceutically acceptable derivative thereof.

20

9. A compound as claimed in any one of claims 1 to 7 for use in therapy.


10. The use of a compound as claimed in any one of claims 1 to 7 in the manufacture of a medicament for use in the treatment or prophylaxis of systemic or topical microbial infections in a human or animal body.

25

11. The use of a compound as claimed in any one of claims 1 to 7 for use in the treatment or prophylaxis of systemic or topical microbial infections in a human or animal body.


30

12. A method for the treatment of the human or non-human animal body to combat microbial infection comprising administration to a body in need of such treatment of an effective amount of a compound as claimed in any one of claims 1 to 7.
13. A pharmaceutical composition comprising at least one compound as claimed in any one of claims 1 to 7 in association with a pharmaceutically acceptable excipient, diluent and/or carrier.
14. A compound of formula (IA)



wherein

- 15 A is a bivalent radical selected from $-C(O)-$, $-C(O)NH-$, $-NHC(O)-$, $-N(R^7)-CH_2-$, $-CH_2-N(R^7)-$, $-CH(NR^8R^9)-$ and $-C(=NR^{10})-$;
- R^1 is $-O(CH_2)_dXR^{11}$;
- R^2 is hydrogen or a hydroxyl protecting group;
- R^3 is hydrogen, C_{1-4} alkyl, or C_{3-6} alkenyl optionally substituted by 9 to 10 membered fused bicyclic heteroaryl;
- 20 R^4 is hydroxy, C_{3-6} alkenyloxy optionally substituted by 9 to 10 membered fused bicyclic heteroaryl, or C_{1-6} alkoxy optionally substituted by C_{1-6} alkoxy or $-O(CH_2)_eNR^7R^{12}$,
- R^5 is hydroxy, or
- R^4 and R^5 taken together with the intervening atoms form a cyclic group having the
- 25 following structure:



wherein Y is a bivalent radical selected from $-CH_2-$, $-CH(CN)-$, $-O-$, $-N(R^{13})-$ and $-CH(SR^{13})-$;

R⁶ is hydrogen or fluorine;


R⁷ is hydrogen or C₁₋₆alkyl;

R⁸ and R⁹ are each independently hydrogen, C₁₋₆alkyl, -C(=NR¹⁰)NR¹⁴R¹⁵ or -C(O)R¹⁴, or


5 R⁸ and R⁹ together form =CH(CR¹⁴R¹⁵)_faryl, =CH(CR¹⁴R¹⁵)_fheterocyclyl, =CR¹⁴R¹⁵ or =C(R¹⁴)C(O)OR¹⁴, wherein the alkyl, aryl and heterocyclyl groups are optionally substituted by up to three groups independently selected from R¹⁶;

R¹⁰ is -OR¹⁷, C₁₋₆alkyl, -(CH₂)_garyl, -(CH₂)_gheterocyclyl or -(CH₂)_hO(CH₂)_iOR⁷,
 10 wherein each R¹⁰ group is optionally substituted by up to three groups independently selected from R¹⁶;

R¹¹ is a heterocyclic group having the following structure:



15 or



R¹² is hydrogen or C₁₋₆alkyl;

20 R¹³ is hydrogen or C₁₋₄alkyl substituted by a group selected from optionally substituted phenyl, optionally substituted 5 or 6 membered heteroaryl and optionally substituted 9 to 10 membered fused bicyclic heteroaryl;

R¹⁴ and R¹⁵ are each independently hydrogen or C₁₋₆alkyl;

25 R¹⁶ is halogen, cyano, nitro, trifluoromethyl, azido, -C(O)R²¹, -C(O)OR²¹, -OC(O)R²¹, -OC(O)OR²¹, -NR²²C(O)R²³, -C(O)NR²²R²³, -NR²²R²³, hydroxy, C₁₋₆alkyl, -S(O)_kC₁₋₆alkyl, C₁₋₆alkoxy, -(CH₂)_maryl or -(CH₂)_mheteroaryl, wherein the alkoxy group is optionally substituted by up to three groups independently selected from -NR¹⁴R¹⁵, halogen and -OR¹⁴, and the aryl and heteroaryl groups are optionally substituted by up to five groups independently selected from halogen, cyano, nitro, trifluoromethyl, azido, -C(O)R²⁴, -C(O)OR²⁴, -OC(O)OR²⁴, -NR²⁵C(O)R²⁶, -C(O)NR²⁵R²⁶, -NR²⁵R²⁶,
 30 hydroxy, C₁₋₆alkyl and C₁₋₆alkoxy;

R¹⁷ is hydrogen, C₁₋₆alkyl, C₃₋₇cycloalkyl, C₃₋₆alkenyl or a 5 or 6 membered heterocyclic group, wherein the alkyl, cycloalkyl, alkenyl and heterocyclic groups are optionally substituted by up to three substituents independently selected from optionally

substituted 5 or 6 membered heterocyclic group, optionally substituted 5 or 6 membered heteroaryl, $-\text{OR}^{27}$, $-\text{S}(\text{O})_n\text{R}^{27}$, $-\text{NR}^{27}\text{R}^{28}$, $-\text{CONR}^{27}\text{R}^{28}$, halogen and cyano;

R^{18} is hydrogen, $-\text{C}(\text{O})\text{OR}^{29}$, $-\text{C}(\text{O})\text{NHR}^{29}$ or $-\text{C}(\text{O})\text{CH}_2\text{NO}_2$;

R^{19} is hydrogen, C_{1-4} alkyl optionally substituted by hydroxy or C_{1-4} alkoxy, C_{3-7} cycloalkyl, or optionally substituted phenyl or benzyl;

R^{20} is halogen, C_{1-4} alkyl, C_{1-4} thioalkyl, C_{1-4} alkoxy, $-\text{NH}_2$, $-\text{NH}(\text{C}_{1-4}\text{alkyl})$ or $-\text{N}(\text{C}_{1-4}\text{alkyl})_2$;

R^{21} is hydrogen, C_{1-10} alkyl, $-(\text{CH}_2)_p\text{aryl}$ or $-(\text{CH}_2)_p\text{heteroaryl}$;

R^{22} and R^{23} are each independently hydrogen, $-\text{OR}^{14}$, C_{1-6} alkyl, $-(\text{CH}_2)_q\text{aryl}$ or $-(\text{CH}_2)_q\text{heterocyclyl}$;

R^{24} is hydrogen, C_{1-10} alkyl, $-(\text{CH}_2)_r\text{aryl}$ or $-(\text{CH}_2)_r\text{heteroaryl}$;

R^{25} and R^{26} are each independently hydrogen, $-\text{OR}^{14}$, C_{1-6} alkyl, $-(\text{CH}_2)_s\text{aryl}$ or $-(\text{CH}_2)_s\text{heterocyclyl}$;

R^{27} and R^{28} are each independently hydrogen, C_{1-4} alkyl or C_{1-4} alkoxy C_{1-4} alkyl;

R^{29} is hydrogen or C_{1-6} alkyl optionally substituted by up to three groups independently selected from halogen, C_{1-4} alkoxy, $-\text{OC}(\text{O})\text{C}_{1-6}$ alkyl and $-\text{OC}(\text{O})\text{OC}_{1-6}$ alkyl;


R^{30} is hydrogen, C_{1-4} alkyl, C_{3-7} cycloalkyl, optionally substituted phenyl or benzyl, acetyl or benzoyl;

R^{31} is hydrogen or R^{20} , or R^{31} and R^{19} are linked to form the bivalent radical $-\text{O}(\text{CH}_2)_2-$ or $-(\text{CH}_2)_t-$;

X is $-\text{U}(\text{CH}_2)_v\text{B}-$, $-\text{U}(\text{CH}_2)_v-$ or a group selected from:



and



25

U and B are independently a divalent radical selected from $-\text{N}(\text{R}^{30})-$, $-\text{O}-$, $-\text{S}(\text{O})_z-$, $-\text{N}(\text{R}^{30})\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{N}(\text{R}^{30})-$ and $-\text{N}[\text{C}(\text{O})\text{R}^{30}]_2-$;

W is $-\text{C}(\text{R}^{31})-$ or a nitrogen atom;

30 d is an integer from 2 to 6;

e is an integer from 2 to 4;

f, g, h, m, p, q, r and s are each independently integers from 0 to 4;

i is an integer from 1 to 6;

j, k, n and z are each independently integers from 0 to 2;

35 t is 2 or 3;

v is an integer from 2 to 8;

or a pharmaceutically acceptable derivative thereof.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

BEST AVAILABLE COPY